

Metal biouptake by actively growing cells of metal-tolerant bacterial strains

Authors: Ganiyu Oladunjoye Oyetibo, Matthew Olusoji Ilori, Oluwafemi Sunday Obayori, Olukayode Oladipo Amund

Abstract

Metal uptake potentials of *Pseudomonas aeruginosa* CA207Ni, *Burkholderia cepacia* CA96Co, *Rhodococcus* sp. ALO3Ni, and *Corynebacterium kutscheri* FL108Hg were studied to determine their competence in detoxification of toxic metals during growth. Metabolism-dependent metal biouptake of the bacteria revealed appreciable uptake of the metals (57–61, 10–30, 23–60, and 10–16 mg g dw⁻¹ of Ni²⁺, Cr⁶⁺, Co²⁺, and Cd²⁺, respectively) from medium, after initial drop in pH, without lag phase. The bacteria exhibited 95–100 % removal efficiency for the metals from aqueous medium as 21 (±0.8)–84 (±2.0) concentration factors of the metals were transported into the bacterial systems. Passive adsorption onto the cell surfaces occurred within 2-h contact, and afterwards, there was continuous accumulation for 12 days. Biosorption data of the bacteria were only fitted into Langmuir isotherm model when strains AL96Co, CA207Ni, and ALO3Ni interacted with Ni²⁺, achieving maximum uptake of 9.87, 2.72, and 2.69 mg g dw⁻¹, respectively. This study established that the actively growing bacterial strains displayed, at least, 97.0 % (±1.5) continuous active removals of metals upon adsorption. The bacteria would be good candidates for designing bioreactor useful in the detoxification campaign of heavy metal-polluted systems.

Keywords: Biouptake, Heavy metal, Bioaccumulation, Bacteria, Detoxification

DOI: <https://doi.org/10.1007/s10661-015-4731-z>

Journal of Environmental Monitoring and Assessment

Published by: Springer International Publishing, on 2015/8/1