

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/333353361>

Reaction Kinetics for Nitrosation of DAF-2 in Air Saturated Nitric Oxide Solution

Article in *Nature and Science* · January 2012

CITATION

1

READS

70

4 authors, including:



Emmanuel Olasehinde

Federal University of Technology, Akure

28 PUBLICATIONS 189 CITATIONS

[SEE PROFILE](#)



Emmanuel Abata

Federal University of Technology, Akure

17 PUBLICATIONS 16 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Production and Characterization of Green Inhibitors for Industrial Use [View project](#)

Reaction Kinetics for Nitrosation of DAF-2 in Air Saturated Nitric Oxide Solution

Olasehinde, E. F.^{1,2}, Ogunsuyi, H.O.², Abata, E. O.² Sakugawa, H.¹

¹ Graduate School of Biosphere Science, Hiroshima University, Japan

² Department of Chemistry, Federal University of Technology, Akure, Nigeria

folasemb2004@yahoo.com

Abstract: Understanding the reaction rate constant of a probe compound with its target molecule is essential for its selection and use in biological and non-biological systems. Over the past decade, the probe, 4, 5-diaminofluorescein (DAF-2) has been widely employed for the detection and imaging of nitric oxide (NO•) in various media. However, the rate constant for the nitrosation of DAF-2 in air-saturated nitric oxide solution is far from being understood. For the first time, we have determined the second order rate constant for the reaction of DAF-2 with NO• (k_2) in air-saturated solution using competition kinetics method. An alternative competition kinetics method which involves a reciprocal plot of the reactant (DAF-2) and triazolofluorescein (DAF-2T), the product formed from the reaction of DAF-2 with NO• was developed and compared to the standard competition kinetics method. Our particular approach in this system is based on the use of oxy-hemoglobin (HbO₂), a potent scavenger of NO• against the DAF-2 which serves as the detector molecule. The product, DAF-2T, is separated from the reaction mixture by reversed-phase high performance liquid chromatography and its fluorescence intensity signals were measured at excitation and emission of 495 and 515 nm respectively. The results showed that the second order reaction rate constant of DAF-2 with NO• in air-saturated aqueous solution are comparable, with the average value of $(6.28 \pm 0.45) \times 10^6 \text{ M}^{-1}\text{s}^{-1}$. Also, DAF-2 can react with NO• directly thereby by-passing the N₂O₃-mediated formation of DAF-2T at low NO• formation rates.

[Olasehinde, EF, Ogunsuyi, HO, Abata, OE, Sakugawa, H. **Reaction Kinetics for Nitrosation of DAF-2 in Air Saturated Nitric Oxide Solution.** *Nat Sci* 2012;10 (11):129-136]. (ISSN: 1545-0740).

<http://www.sciencepub.net/nature>. 19

Keywords: Rate constant; Competition kinetics; Nitric oxide; DAF-2; DAF-2T; HPLC



1. Introduction

Fluorescein is widely used in biology as a fluorophore because of its convenient absorption wavelength for biological measurement, high extinction coefficient and high fluorescence quantum yield in water (Katayama et al., 1998; Kojima et al., 1998 a, 1998 b; Kojima and Nagano, 2000; Nagano, 1999). Over the past decade, DAF-2 has been extensively used as a chemical probe for the detection and imaging of NO• generated under physiological conditions or nitrite ion (NO₂⁻) under acidic conditions (Leikert et al., 2001; Nakatsubo et al., 1998; Planchet and Kaiser, 2006; Rathel et al., 2003; Suzuki et al., 2002). In fact, it has been reported that diaminofluoresceins (DAFs) have been applied to the determination of NO• in more than 100 scientific reports in the last 2 years (Gomes et al., 2006). Recently, we reported for the first time the application of this probe to the measurement of photochemically generated NO• in natural waters (Olasehinde et al., 2009; Olasehinde et al., 2010). However, despite the compelling evidence that DAF-2 can be used for the sensitive analysis of NO• in various media, the reaction rate constant of this probe with NO• has always been a puzzle. It should be pointed out that the reactive oxygen species along with other free radicals can react with

NO• in competition with reaction with the probe (Olasehinde et al., 2009; Olasehinde et al., 2010; Zafiriou and Mc Farland, 1980). Therefore, information on the reaction rate constant of DAF-2 with NO• in air-saturated solution is critical to assessing the competition quantitatively, comparing the products of rate constant and the concentration. This will ultimately unravel the potency of DAF-2 for nitric oxide assay.

Pulse radiolysis is a direct method and is known to be the best for rate constant determinations (Ashton et al., 1995; Joseph and Aravindakumar, 2000). However, its use is often limited by its expensive nature and hence is not available to many radical chemists. Therefore, the competition kinetics method is employed for rate constant determinations (Gholami et al 1998; Kochany and Bolton, 1991; Manoj and Aravindakumar, 2003; Onstein et al., 1999). On this basis, we have carried out a series of experiments to determine second order reaction rate constant of DAF-2 with NO• (k_2) in air-saturated aqueous solution using competition kinetics methods. In the competition assay, information on the reaction rate can be obtained when the chemical reaction between the target molecule and a potential scavenger is examined in the presence of a detector molecule.

In this study, the detector molecule is DAF-2 and HbO₂ serves as the scavenger. Nitric oxide was generated *in situ* according to the equation (1) by the irradiation of NO₂⁻ with simulated light in air-saturated aqueous solution, which reacted competitively with the scavenger oxy-hemoglobin (HbO₂) and the detector molecule (DAF-2). It is well known that all amine-based probes such as DAF-2 are unreactive toward NO• without initial oxidation (Jourdeuil, 2002; Espey et al., 2002; Williams, 1997; Wardman, 2007). Jourdeuil (2002) suggested that •OH and NO₂[•] are capable of rapid reaction with DAF-2 to generate DAF-2 radical intermediate, which can then react with NO• to form DAF-2T. As pointed out by Wardman (2007), DAF-2 radical intermediate can also be generated from the photoexcitation of DAF-2 probe. In our system, a possible mechanism for the formation of DAF-2T involves a one-electron oxidation of DAF-2 via the reaction of DAF-2 with free radicals (such as hydroxyl radicals) or/and photosensitized oxidation of DAF-2 to a free radical derivative, which reacts with NO• via radical-radical combination (eqs 2 and 3; Williams, 1997; Wardman, 2007).



The product, DAF-2T, which is formed in the reaction between NO• and DAF-2 (Figure 1) in air-saturated solution is quantified by reversed-phase HPLC equipped with a fluorescence detector. The decrease in the amount of DAF-2T that was formed due to the presence of HbO₂ depends on the initial concentrations of DAF-2 and HbO₂, the concentration of NO• generated from NO₂⁻ and the ratio of the rate constants of (3) and (4). Since the reaction rate constant of HbO₂ with NO• is known, the value for (k₂) was determined using different ratios of HbO₂ and DAF-2.




Fig.1 Structures of DAF-2 and DAF-2T

2. Material and Methods

2.1. Reagents and Chemicals

All reagents were reagent grade and used as received unless otherwise stated. All solutions were prepared with ultra-pure water obtained from a Milli-Q Plus system (Millipore; $\geq 18.2 \text{ M}\Omega \text{ cm}$). Acetonitrile and benzene (HPLC grade, > 99.5), and disodium hydrogen phosphate heptahydrate (guaranteed reagent) were obtained from Nacalai Tesque, (Tokyo, Japan). Sodium dithionite, sodium azide, bovine hemoglobin and sodium phosphate monobasic monohydrate were obtained from Sigma - Aldrich Japan. DAF-2 was obtained from Daiichi Pure Chemicals (Japan) and DAF-2T from Alex Biochemicals (Japan). NO₂⁻ standard solution (1000 mg L⁻¹) was obtained from Kanto Chemical Co. Inc.

2.2. Equipment

An HPLC system consisting of a PU-2089 plus pump (Jasco, Japan), a Rheodyne injection valve (Cotati, CA, USA) with a 100 μ L sample loop and a FP-2020 plus intelligent fluorescence detector (Jasco, Japan) interfaced with a C-R6A Chromatopac integrator (Shimadzu, Japan) was used. The separations were carried out on a RP-18 GP column (150 \times 4.6mm I.D., 5 μ m) from Kanto Kagaku (Japan) with 10 mM phosphate buffer saline (PBS) at pH 7.4 with 6% acetonitrile as eluent at a flow rate of 1 mL min⁻¹. The detector was operated at 495nm and 515 nm for excitation and emission respectively.

2.3. Purification of HbO₂

To prepare HbO₂ from the commercial bovine hemoglobin, 1 mM sodium dithionite was added to a solution of bovine hemoglobin in 100 mM potassium phosphate at pH 7.0 to reduce any met-hemoglobin present in the solution. The mixture was dialyzed overnight at 4°C in 100 mM potassium phosphate, pH 7.0, to remove the residual dithionite. The concentration was determined by the previously reported method (Herold and Rock, 2003).

2.4. Irradiation Experiment

A solar simulator (Oriel model 81160-1000, Oriel corp.) equipped with a 300 W Xe lamp (ozone free, model 6258, Oriel corp.) was employed for the irradiation experiments. To simulate actual solar irradiance, wavelengths less than 300 nm were filtered out by optical filters (Oriel AM 0 and AM1.0, Oriel Instruments). The daily actinic flux of the solar simulator was determined by chemical actinometry (2-Nitrobenzaldehyde, 2-NB) (Arakaki et al., 1999). The photochemical reactions were performed in a custom-made quartz cuvette cell (Workshop for advanced Techniques, Hiroshima University). In this system, samples were illuminated in a 7-mL quartz cuvette cell. The solution inside the cell was gently stirred with a Teflon stirring bar and maintained at ca. 20°C using a Neslab RTE 111 recirculatory water bath. Aliquots of the irradiated

samples were removed at suitable intervals for HPLC analysis.

3. Results and Discussion

3.1 System Validation

3.1.1 Degradation of DAF-2 and Formation of DAF-2T

In our previous study (Olasehinde et al., 2009), we reported the selectivity towards $\text{NO}\bullet$ of the reaction transforming DAF-2 into DAF-2T. It was shown that the signal intensity of DAF-2T increased with an increase in the NO_2^- concentration. This suggests that DAF-2 reacts with the photoformed $\text{NO}\bullet$ generated from NO_2^- photolysis to produce DAF-2T in a concentration-dependent manner.

In this study, we assumed that DAF-2 concentration remained essentially constant during the course of the reaction and that DAF-2T formation was negligible. These assumptions were checked by irradiating 5 mL of Milli-Q water containing 5 μM DAF-2 for 30 min. An aliquot of the reaction mixture was subjected to HPLC analysis before and after the illumination. Concentrations of DAF-2 and DAF-2T were determined using calibration standards prepared in phosphate buffer (0.1 M, pH 7.4) and run on the day of the illumination experiment. The analysis of these samples ($n = 3$) by HPLC showed a $< 2\%$ reduction in the concentration of each DAF-2. Meanwhile, DAF-2T was detectable in small amounts, but the concentration was always similar in the pairs of samples. As reported by Olasehinde et al (2009), DAF-2T was present as an impurity in the DAF-2, as opposed to being formed during the reaction period. We and others have also shown that stable oxidized forms of $\text{NO}\bullet$, such as NO_2^- and NO_3^- , and other reactive oxygen species, such as $\bullet\text{OH}$, $\text{O}_2\bullet^-$, H_2O_2 , and ONOO^- , do not react with DAF-2 to yield any fluorescent product and that DAF-2T is not formed in the absence of $\text{NO}\bullet$ (Gomes et al., 2006; Olasehinde et al., 2009; Kojima et al., 1998 b). This clearly indicates that the increase in fluorescence intensity of DAF-2T was dependent on the $\text{NO}\bullet$ concentration rather than the OH radicals in equation (1).

However, it is a well-known fact that the reactions of $\bullet\text{OH}$ and $\text{NO}\bullet$ are diffusion-controlled (Mack and Bolton, 1999) with a second-order rate constant of $1.0 \times 10^{10} \text{ M}^{-1}\text{s}^{-1}$. To examine the scavenging effects of OH radicals on $\text{NO}\bullet$ in equation 1, a 5 mL Milli-Q water containing 10 μM NO_2^- was irradiated with 5 μM DAF-2 in the absence and presence of 1.2 mM benzene. We experimentally confirmed that 1.2 mM benzene is sufficient to effectively scavenge all $\bullet\text{OH}$ formed from the irradiation of 10 μM NO_2^- in Milli-Q water (data not shown), which agrees with the conclusion of the previous investigator (Nakatani et al., 2007). If the

scavenging effect of $\bullet\text{OH}$ on the photoformed $\text{NO}\bullet$ were not negligible, the fluorescence intensity of DAF-2T would be expected to increase significantly upon the addition of $\bullet\text{OH}$ scavenger to compensate for the loss of $\text{NO}\bullet$ due to its reaction with $\bullet\text{OH}$.

Contrarily, the result showed no appreciable difference between the fluorescence intensity of DAF-2T formed in the presence and absence of benzene (Figure. 2a), suggesting the negligible effect of OH radicals on the nitric oxide generated in equation 1. Further, it has been shown that 2 μM DAF-2 was sufficient to effectively scavenge all $\text{NO}\bullet$ formed from the irradiation of 10 μM NO_2^- in Milli-Q water in the presence of other *in situ* generated radicals (Olasehinde et al., 2009). In this study, 5 μM DAF-2 was employed to scavenge all $\text{NO}\bullet$ formed in the presence of other *in situ* generated scavengers.




Figure 2a: The signal intensity of DAF-2T produced by the irradiation of 10 μM NO_2^- and 5 μM DAF-2 for 20 min in air-saturated solution (control) and with the addition of 1.2 mM benzene. Each experiment was conducted in triplicate.



If we assume that the steady-state concentration of $\bullet\text{OH}$ in equation 1 is as high as $\sim 10^{-15} \text{ M}$ (a reasonable upper limit based on the $\bullet\text{OH}$ photoformation rates from NO_2^- photolysis and the scavenging rate constant of $\bullet\text{OH}$ by NO_2^- , ignoring loss of $\bullet\text{OH}$ due to other mechanisms) and the second order rate constant for the recombination reaction of the $\text{NO}\bullet$ and OH radicals of $1.0 \times 10^{10} \text{ M}^{-1}\text{s}^{-1}$ (Mack and Bolton, 1999; Buxton et al., 1988), then equation 5 proceeds with a first-order rate constant of $1.0 \times 10^5 \text{ s}^{-1}$.



In contrast, if k_2 is $6.28 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ and the DAF-2 concentration is $5 \mu\text{M}$, equation (3) has a first-order rate constant of 31 s^{-1} , a 6-order of magnitude higher than equation (5). From this simple analogy, it is obvious that the effect of $\bullet\text{OH}$ on $\text{NO}\bullet$ in equation 1 is negligible at the optimum DAF-2 concentration employed in this study.

3.1.2 Specificity of DAF-2 for NO

The nitrosating species in the reaction of DAF-2 with $\text{NO}\bullet$ in air-saturated solution has long been a subject of controversy. A number of studies have established that the nitrosating species in the reaction between DAF-2 and $\text{NO}\bullet$ in air-saturated solution is N_2O_3 , suggesting that $\text{NO}\bullet$ does not nitrosate DAF-2 directly and that the nitrosation of DAF-2 in aerobic solution is consistent with the formation of N_2O_3 consequent to $\text{NO}\bullet$ autoxidation (Gomes et al., 2006; Nakastubo et al. 1998; Rathel et al., 2003; Nagano et al., 1995; Leikert et al., 2001, Kojima et al., 1998b; Nakastubo et al. 1998; Rathel et al., 2003; Nagano et al., 1995; Leikert et al., 2001, Kojima et al, 1998b).



However, recent studies have shown that DAF-2T formation is not strictly related to the reaction with N_2O_3 (Espey et al, 2002; Jourdeheu et al., 2002). These authors hypothesized that the triazole product (DAF-2T) may be formed predominantly by oxidative nitrosylation, a process that involves oxidation of the probe followed by direct reaction with $\text{NO}\bullet$ (Espey et al., 2002). In a separate study using 2, 3-diaminonaphthalene (DAN), an analogue of DAF-2, for trapping $\text{NO}\bullet$ in seawater, the results showed that the fluorescence intensity between oxygenated and de-oxygenated DAN solutions was in the range of detection error, suggesting that DAN can react with $\text{NO}\bullet$ directly in aerobic solutions (Liu et al., 2009).

To better understand the contribution of N_2O_3 to DAF-2T fluorescence, we performed a series of inhibition experiments by irradiating 5 mL of Milli-Q water containing air-saturated solution of $10 \mu\text{M}$ NO_2^- and $5 \mu\text{M}$ DAF-2 in the presence and absence of N_2O_3 scavenger sodium azide. Although the rate constant for the reaction of DAF-2 with N_2O_3 (k_8) is not known, azide reacts with N_2O_3 at a

diffusion controlled rate, with a second order reaction rate constant of $2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ (Williams, 1997). If we assume that k_8 is as high as $\sim 10^9 \text{ M}^{-1} \text{ s}^{-1}$, and that N_2O_3 were the intermediate responsible for the nitrosation of DAF-2 in air-saturated solution, equation 8 proceeds with a first-order rate constant of $5.0 \times 10^3 \text{ s}^{-1}$ at $5 \mu\text{M}$ DAF-2 employed in this study. However, using the published rate constant of N_2O_3 with azide (Williams, 1997), and at 0.5 mM $[\text{N}_3^-]$, about 99.5 % of the observed DAF-2T should be quenched. In contrast, addition of 0.5 mM sodium azide showed no detectable decrease in the fluorescence intensity of DAF-2T (Figure 2b), a discrepancy which appears to preclude the mechanisms described by eqs 6-8 under our experimental conditions.




Figure 2b: The signal intensity of DAF-2T generated by the irradiation of $10 \mu\text{M}$ NO_2^- with $5 \mu\text{M}$ DAF-2 for 20 min in air-saturated solution (control), with the addition of 0.5 mM sodium azide, and when the solution was purged with N_2 for 30 min prior to irradiation. Each experiment was conducted in triplicate.

As shown in Figure 2c, when $\text{NO}\bullet$ scavenger HbO_2 (0.1 - $0.4 \mu\text{M}$) was added to the reaction mixture containing $10 \mu\text{M}$ NO_2^- with $5 \mu\text{M}$ DAF-2 followed by irradiation, a concentration-dependent reduction in the fluorescence intensity of DAF-2T was observed. Furthermore, the possibility of DAF-2T formation in anaerobic solution, bypassing the requirement for $\text{NO}\bullet$ autoxidation for the formation of DAF-2T was investigated. In each of these experiments, a 5-mL Milli-Q water containing $10 \mu\text{M}$ NO_2^- and $5 \mu\text{M}$ DAF-2 was purged with a high-purity N_2 gas for 30 min to remove oxygen prior to irradiation. The solutions were then irradiated for

20 min and analyzed for DAF-2T. The difference in the fluorescence intensity of DAF-2T in the aerobic solutions compared to the anaerobic solutions was < 5%, which is within the analytical error (See Figure 2b).




Figure 2c Signal intensities of DAF-2T produced by the irradiation of 10 μM NO_2^- and 0.5 μM DAF-2 in Milli-Q water with and without HbO_2 as a function of irradiation time. Closed diamonds: 0 μM HbO_2 ; closed rectangles: 0.1 μM HbO_2 ; open rectangles: 0.2 μM HbO_2 ; closed circles: 0.3 μM HbO_2 ; open circles: 0.4 μM HbO_2 .

To further clarify the likelihood of equation 6 to generate N_2O_3 compared to the loss of $\text{NO}\cdot$ via equation 3, the first order rate constants for eqs 6 and 3 were estimated. If we assume that the steady-state concentration of $\text{NO}\cdot = \sim 10^{-15}$ M (based on the ratio of the $\text{NO}\cdot$ photo-formation rate from NO_2^- and the scavenging rate by NO_2^- , ignoring the loss of $\text{NO}\cdot$ due to other processes), $k_6 = 1.1 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$ (Mack and Bolton, 1999), the forward half-reaction for equation 6 proceeds with a first-order rate constant of $1.1 \times 10^{-6} \text{ s}^{-1}$. As mentioned earlier, equation 3 proceeds with a first-order rate constant of 31.1 s^{-1} indicating a 7-order of magnitude higher than the equation 6.

Taken together, these data signify that the nitrosating species under our experimental conditions is $\text{NO}\cdot$ rather than the addition of N_2O_3 . This is reasonable because previous reports (Espey et al., 2002; Wink et al., 1994), demonstrated that at low $\text{NO}\cdot$ formation rates ($\sim \text{nM NO}\cdot / \text{min}$), the half-life of NO in aerobic environments is potentially quite long, which may allow other pathways such as the reaction of $\text{NO}\cdot$ with oxy-hemoglobin (or other $\text{NO}\cdot$ probe compounds) to predominate over the nitrosation addition by N_2O_3 . This result suggests that DAF-2 can react with $\text{NO}\cdot$ directly (as in equation 3), thereby by-passing the N_2O_3 -mediated formation of DAF-2T. However, nitrosation via $\text{NO}\cdot$ autoxidation may be the dominant route when relatively high rates

of $\text{NO}\cdot$ formation occur under aerobic condition which would favour nitrosation through N_2O_3 formation subsequent to the reaction of NO_2^- with $\text{NO}\cdot$ (Espey et al., 2002).

3.2 Determination of K_2 using Standard Competition Kinetics

We employed the standard competition kinetics method (Buxton et al., 1988) to determine the reaction rate constant of DAF-2 with $\text{NO}\cdot$ (k_2) in air-saturated aqueous solution. Nitric oxide radical was generated by irradiating 10 μM NO_2^- in a 7-mL quartz cell. DAF-2 reacts with $\text{NO}\cdot$ in air-saturated solution yielding the highly fluorescent DAF-2T. Also, HbO_2 reacts with $\text{NO}\cdot$ to produce methemoglobin (metHb), with a second order reaction rate of $3.7 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ (Gross and Lane, 1999). In experimental solution containing DAF-2 without any other $\text{NO}\cdot$ scavenger, all $\text{NO}\cdot$ formed reacts with DAF-2 to produce DAF-2T. The total amount of DAF-2T formed under this condition is denoted as Q_T . However, if HbO_2 is present in the experimental solution with DAF-2, both reactants compete for the $\text{NO}\cdot$ produced which leads to the attenuation of the signal intensity of DAF-2T compared to the reaction of DAF-2 with $\text{NO}\cdot$ in the absence of HbO_2 . In this system, the amount of DAF-2T formed is a fraction of the total $\text{NO}\cdot$ generated which is denoted as Q . If the concentration of DAF-2T is represented by Q and Q' represents the concentration of metHb, therefore the total scavengable $\text{NO}\cdot$ under this condition is given as:

$$Q_T = Q + Q' \quad (9)$$

$$(dQ/dt) / (dQ'/dt) = (Q / Q')$$

$$(dQ/dt) = k_2 [\text{NO}\cdot] [\text{DAF-2}]$$

$$(dQ'/dt) = k_3 [\text{NO}\cdot] [\text{HbO}_2]$$

$$Q / Q' = (k_2 [\text{NO}\cdot] [\text{DAF-2}]) / (k_3 [\text{NO}\cdot] [\text{HbO}_2]) \quad (10)$$

Since Q is the measurable product and Q' is expressed in terms of Q

Therefore

$$Q' = Q_T - Q \quad (11)$$

Substitute eq 11 in eq 10

$$Q / (Q_T - Q) = (k_2 [\text{DAF-2}]) / (k_3 [\text{HbO}_2]) \quad (12)$$

Inverse of eq 12 gives

$$(Q_T - Q) / Q = (k_3 [\text{HbO}_2]) / (k_2 [\text{DAF-2}]) \quad (13)$$

eq 13 is simplified as

$$\{Q_T / Q\} = 1 + \{(k_3 [\text{HbO}_2]) / (k_2 [\text{DAF-2}])\} \quad (14)$$

Let

$$Q_T = [\text{DAF-2T}]_0, Q = [\text{DAF-2T}]'$$

Therefore, eq 14 becomes

$$[\text{DAF-2T}]_0 / [\text{DAF-2T}]' = 1 + \{(k_3 [\text{HbO}_2]) / (k_2 [\text{DAF-2}])\} \quad (15)$$

In equation 15, $[\text{DAF-2T}]'$ and $[\text{DAF-2T}]_0$ are the concentrations of DAF-2T produced from the reaction of DAF-2 and $\text{NO}\cdot$ in air-saturated solutions with and without HbO_2 respectively, $[\text{HbO}_2]$ is the initial concentration of HbO_2 , $[\text{DAF-2}]$ is the initial

concentration of DAF-2, k_3 is the reaction rate constant of HbO_2 with $\text{NO}\cdot$ ($\text{M}^{-1}\text{s}^{-1}$) and k_2 is the reaction rate constant of DAF-2 with $\text{NO}\cdot$ ($\text{M}^{-1}\text{s}^{-1}$). We carried out a series of experiments by 30-min irradiation of $10\ \mu\text{M}$ NO_2^- with (0.05 – 0.1 μM) HbO_2 varying the concentration of DAF-2 from (0.1 - 0.5 μM). When the DAF-2 concentration was held constant and the concentration of HbO_2 was increased, a concentration-dependent reduction in the amount of DAF-2T ' was observed. Contrarily, when the concentration of HbO_2 was held constant while increasing the concentration of DAF-2, the signal intensity of DAF-2T ' was enhanced to compensate for the increase in the concentration of DAF-2. However, to determine the concentration of the Q_T produced $10\ \mu\text{M}$ NO_2^- was irradiated with $5\ \mu\text{M}$ DAF-2 in the absence of HbO_2 for 30 min. As reported earlier (Olasehinde et al., 2009), the concentration dependence of DAF-2 on $\text{NO}\cdot$ generated from the irradiation of $10\ \mu\text{M}$ NO_2^- in Milli-Q water revealed that $5\ \mu\text{M}$ DAF-2 is sufficient to effectively scavenge all $\text{NO}\cdot$ formed. A plot of $[\text{DAF-2T}]_0 / [\text{DAF-2T}]'$ vs $[\text{HbO}_2] / [\text{DAF-2}]$ for separate experiments with solutions of varying $[\text{HbO}_2] / [\text{DAF-2}]$ ratio, based on eq 15 is shown in Figure 3. Thus, the value for k_2 estimated from the slope of this graph and eqn 15 was $(6.17 \pm 0.26) \times 10^6\ \text{M}^{-1}\ \text{s}^{-1}$ (mean \pm standard deviation, $n=3$).




Figure 3: Graph for the calculation of the rate constant for the reaction of DAF-2 with $\text{NO}\cdot$ radical using the competition kinetics method, correlation coefficient (r^2) = 0.999.

3.3. Determination of k_2 using Alternate Competition Kinetics Method

We proposed an alternate competition kinetics method for the determination of reaction rate constant of DAF-2 with $\text{NO}\cdot$ to ensure that the result obtained from the standard competition kinetics method was not a mere artifact. The following mathematical derivation will aid in the determination of k_2 .

$$\text{Given that } R_{\text{NO}} = R_{\text{DAF-2T}} / (F_{\text{DAF-2, NO}}) (Y_{\text{DAF-2T}}) \quad (16)$$

$$\text{and } F_{\text{DAF, NO}} = (k_2) [\text{DAF-2}] / \sum (k_s, \text{NO} [s]) \quad (17)$$

Where $R_{\text{DAF-2T}}$ is the formation rate of DAF-2T (M/s), $Y_{\text{DAF-2T}}$ is the yield of DAF-2T per DAF-2

oxidized by $\text{NO}\cdot$, $F_{\text{DAF-2, NO}}$ is the fraction of DAF-2 that reacts with $\text{NO}\cdot$, k_2 is the reaction rate constant of DAF-2 with $\text{NO}\cdot$ ($\text{M}^{-1}\text{s}^{-1}$), R_{NO} is the formation rate of $\text{NO}\cdot$ and $\sum (k_s, \text{NO} [s])$ is the summation of scavenging rate of $\text{NO}\cdot$ with $\text{NO}\cdot$ scavengers in experimental solution.

In an experimental solution containing DAF-2, HbO_2 and other *in situ* generated scavengers of $\text{NO}\cdot$, the summation of scavenging rate of $\text{NO}\cdot$ by the scavengers is given in Eq. 18.

$$\sum (k_s, \text{NO} [s]) = (k_2) [\text{DAF-2}] + k_3 [\text{HbO}_2] + k_a [\text{S}_a] + k_b [\text{S}_b] + \dots + k_n [\text{S}_n] \quad (18)$$

where k_a, \dots, k_n are the rate constants of individual scavengers $\text{S}_a, \dots, \text{S}_n$ with $\text{NO}\cdot$

However, If HbO_2 and DAF-2 are the main scavengers of $\text{NO}\cdot$ in the experimental solution such that the total $\text{NO}\cdot$ consumption excluding HbO_2 and DAF-2 is significantly lower than HbO_2 and DAF-2, that is: $\{ (k_2) [\text{DAF-2}] + k_3 [\text{Hb}] \} \gg \sum (k_a [\text{S}_a] + k_b [\text{S}_b] + \dots + k_n [\text{S}_n])$,

Then

$$\sum (k_s, \text{NO} [s]) = (k_2) [\text{DAF-2}] + k_3 [\text{HbO}_2] \quad (19)$$

Substituting Eq. (19) in Eq. (17) gives

$$F_{\text{DAF-2, NO}} = k_2 [\text{DAF-2}] / \{ k_2 [\text{DAF-2}] + k_3 [\text{HbO}_2] \} \quad (20)$$

Substituting Eq. (20) in Eq. (16) and re-arranged gives

$$R_{\text{DAF-2T}} = R_{\text{NO}} \{ (k_2 [\text{DAF-2}]) (Y_{\text{DAF-2T}}) \} / \{ k_2 [\text{DAF-2}] + k_3 [\text{HbO}_2] \} \quad (21)$$

This can then be solved algebraically in terms of $1/R_{\text{DAF-2T}}$ to give:

$$1/R_{\text{DAF-2T}} = \{ k_2 [\text{DAF-2}] + k_3 [\text{HbO}_2] \} / R_{\text{NO}} \{ (k_2 [\text{DAF-2}]) (Y_{\text{DAF-2T}}) \} \quad (22)$$

Simplifying Eq. (22) gives

$$1/R_{\text{DAF-2T}} = k_3 [\text{HbO}_2] / R_{\text{NO}} \{ (k_2 [\text{DAF-2}]) (Y_{\text{DAF-2T}}) \} + 1/R_{\text{NO}} (Y_{\text{DAF-2T}}) \quad (23)$$

Hence, a plot of $1/R_{\text{DAF-2T}}$ versus $1/[\text{DAF-2}]$ will give a line with

$$\text{slope} = k_3, [\text{Hb}] \times \text{y-intercept} / k_2 \quad (24)$$

Using this methodology, we performed a series of independent experiments by irradiating (5 -10 μM) NO_2^- in Milli-Q water with various concentration of DAF-2 (0.1–1 μM) in the presence of (0.1 – 0.3 μM) HbO_2 . For each experiment, NO_2^- was irradiated with a fixed concentration of HbO_2 while varying the concentration of DAF-2 used. Aliquot of the irradiated sample was withdrawn for analysis at suitable intervals and the concentration of DAF-2T produced was determined. Photo-formation rate of DAF-2T ($R_{\text{DAF-2T}}$) was calculated from the relationship between the cumulative irradiation time and the concentration of DAF-2T. From the relationship between $1/R_{\text{DAF-2T}}$ and $1/[\text{DAF-2}]$ (Figure 4) using Eq. (24), the second order reaction rate constant of DAF-2 with $\text{NO}\cdot$ was estimated to be $(6.39 \pm 0.31) \times 10^6\ \text{M}^{-1}\ \text{s}^{-1}$ (mean \pm standard deviation, $n=3$).




Figure 4. Graph for the calculation of the rate constant for the reaction of DAF-2 with $NO\bullet$ using the alternate competition kinetics method. A plot of reciprocal of DAF-2T photo-formation rate ($1/R_{DAF-2T}$) and inverse of the concentration of added DAF-2 ($1/[DAF-2]$) in Milli-Q water. $[NO_2^-]: 10 \mu M$, $[DAF-2]: (0.1- 0.4 \mu M)$; $[Hb]: 0.1 \mu M$. The values for the y-intercept and slope are $3.45 \times 10^{11} s/M$ and $1.93 \times 10^5 s$ respectively, correlation coefficient (r^2) = 0.993.

4. Conclusions

The second order reaction rate constant of DAF-2 with $NO\bullet$ has been determined. Using the two different competition kinetics approaches, reaction rate constants of DAF-2 with $NO\bullet$ were comparable, with the average value of $(6.28 \pm 0.45) \times 10^6 M^{-1}s^{-1}$. The strong agreement between the two competition kinetics procedures implies that the reaction between DAF-2 and $NO\bullet$ is considerably high. This study also reveals that at low $NO\bullet$ formation rates ($\sim nM NO\bullet / min$), DAF-2 can react with $NO\bullet$ directly thereby bypassing the N_2O_3 -mediated formation of DAF-2T. From the results presented in this study, DAF-2 could serve as a novel probe compound for the sensitive determination of NO in biological samples as well as in aqueous solution.

Acknowledgements

The authors acknowledge with thanks the Japan Society for the Promotion of Science for the financial support through Grants-in-Aid for Scientific Research (B) (18310010).

Corresponding Author:

Dr. Olasehinde Emmanuel F.
Department of Chemistry
Federal University of Technology, Akure
Ondo State, Nigeria
E-mail: folasemb2004@yahoo.com

References

1. Arakaki, T., Miyake, T., Hirakawa, T., Sakagawa, H. pH dependent photo-formation of hydroxyl radical and absorbance of aqueous-phase N(III) (HNO_2 and NO_2). *Environmental Science and Technology*, 1999; 33: 2561–2565.
2. Ashton, L., Buxton, G. V., Stuart, C. R. Temperature dependence of the rate of reaction of OH with some aromatic compounds in aqueous solution. *Chemical Society Faraday Transactions*, 1995; 91: 1631–1633.
3. Buxton, G. V., Greenstock, C. L., Helman, W. P., Ross, A. B. *Journal Physical Chemistry Reference Data*, 1988; 17: 519.
4. Espey, M. G., Thomas, D. D., Miranda, K. M., Wink, D. A. Focusing of nitric oxide mediated nitrosation and oxidative nitrosylation as a consequence of reaction with superoxide. *Proceeding of National Academy of Science, U.S.A.* 2002; 99: 11127–11132.
5. Gholami, M. R., Hoag, C. M., Crawford, C. L., Hanrahan, R. J. An alternate method of treating competition kinetics: reaction of OH radical with $IrBr_6^-$. *Radiation, Physics and Chemistry*, 1998; 53: 505-509.
6. Gomes, A.; Fernandes, E.; Lima, J. L. Use of fluorescence probes for detection of reactive nitrogen species: A review. *Journal of Fluorescence*, 2006; 16: 119–139.
7. Gross, S. S., and Lane, P. Physiological reactions of nitric oxide and hemoglobin: A radical rethink. *Proceeding of the National Academy of Science, USA*, 1999; 96, 9967–9969.
8. Herold, S. and Rock, G. Reaction of Deoxy-, Oxy-, and Methemoglobin with
9. Nitrogen Monoxide: Mechanistic studies of the s-nitrosothiol formation under different mixing conditions. *Journal of Biology and Chemistry*, 2003; 278: 6623 - 6634.
10. Joseph, J. M. and Aravindakumar, C. T. Determination of rate constants for the reaction of hydroxyl radicals with some purines and pyrimidines using sunlight. *Journal of Biochemical and Biophysical Methods*, 2000; 42: 115-124.
11. Jourdeuil, D. Increased nitric oxide-dependent nitrosylation of 4, 5-diaminofluorescein by oxidants: implications for the measurement of intracellular nitric oxide. *Free Radical Biology and Medicine*, 2002; 33: 676 – 680
12. Katayama, Y., Takahashi S., Maeda, M., Design, synthesis and characterization of a novel fluorescent probe for nitric oxide nitrogen monoxide; *Analytica Chimica Acta*, 1998; 365: 159-167.
13. Kochany, J. and Bolton, J. Mechanism of photodegradation of aqueous organic pollutants EPR spin-trapping technique for the determination of OH radical rate constants in the photooxidation

- of chlorophenols following the photolysis of H₂O₂. *Journal of Physical Chemistry*, 1991; 95: 5116–5120.
14. Kojima, H., Nagano, T. Fluorescent indicators for nitric oxide. *Advanced Materials*, 2000; 12: 763-765.
 15. Kojima, H., Nakatsubo, N., Kikuchi, K., Kawahara, S., Kirino, Y., Nagoshi, H., Hirata, Y., Nagano, T. Detection and imaging of nitric oxide with novel fluorescent Indicators: diamino fluoresceins. *Analytical Chemistry*, 1998 b; 70: 2446-2453.
 16. Kojima, H., Sakurai, K., Kikuchi, K., Kawahara, S., Kirino, Y., Nagashi, H., Hirata Y., Nagano, T., Development of a fluorescent indicator for nitric oxide based on the fluorescein chromophore. *Chem. Pharm. Bull.*, 1998 a; 46: 373-375.
 17. Leikert, J. F., Rathel, T. R., Muller, C., Vollmar, A., Dirsch, V. M. Reliable in vitro measurement of nitric oxide released from endothelial cells using low concentration of the fluorescent probe 4,5-diaminofluorescein. *FEBS Letters*, 2001; 506: 131-134.
 18. Liu, C., Zhao, M., Ren, C., Yang, G., Li, P., Han, Y. Direct measurement of nitric oxide in seawater medium by fluorometric method. *Chinese Journal of Analytical Chemistry*, 2009; 37: (10), 1463–1467.
 19. Mack, J. and Bolton, J. R. Photochemistry of nitrite and nitrate in aqueous solution: A review. *Journal of Photochemistry and Photobiology. A: Chem.* 1999; 128, 1-13.
 20. **Manoj, V. M. and Aravindakumar, C. T.** Reaction of hydroxyl radical with s-nitrosylthiols: determination of rate constant and end product analysis **Journal of Organic Bimolecular Chemistry**, 2003; 1: 1171-1175.
 21. Nagano, T. Practical methods for detection of nitric oxide. *Luminescence*, 1999; 14: 283-290.
 22. Nakatani, N., Hashimoto, N., Shindo, H., Yamamoto, M., Kikkawa, M., Sakugawa, H. Determination of photoformation rates and scavenging rate constants of hydroxyl radicals in natural waters using an automatic light irradiation and injection system. *Analytical. Chimica. Acta*, 2007; 581: 260-267.
 23. Nakatsubo, N., Kojima, H., Kikuchi, K., Nagoshi, H., Hirata, Y., Maeda, D., Imai, Y., Irimura, T., Nagano, T. Direct evidence of nitric oxide production from bovine aortic endothelial cells using new fluorescence indicators: diamino fluoresceins. *FEBS Letters*, 1998; 427: 263-266.
 24. Olasehinde, E. F., Takeda, K., Sakugawa, H. Photochemical production and consumption mechanisms of nitric oxide in seawater. *Environmental Science and Technology*, 2010; 44 (22): 8403-8408.
 25. Olasehinde, E. F., Takeda, K., Sakugawa, H. Development of an analytical method for nitric oxide radical determination in natural waters. *Analytical Chemistry*, 2009; 81 (16): 6843–6850.
 26. Onstein, P., Stefan, M., Bolton, J. Competition kinetics method for the determination of rate constants for the reaction of hydroxyl radicals with organic pollutants using the UV/H₂O₂ advanced oxidation technology: The rate constants for the tert-butyl formate ester and 2,4-dinitrophenol. *Journal of Advanced Oxidation Technology*, 1999; 4: 231–236.
 27. Planchet, E. and Kaiser, W. M. Nitric oxide (NO) detection by DAF fluorescein and chemiluminescence: a comparison using abiotic and biotic NO sources. *Journal of Experimental Botany*, 2006; 57: 3043-3055.
 28. Rathel, T. R., Leikert, J. F., Vollmar, A. M., Dirsch, V. M. Application of 4, 5-diaminofluorescein to reliably measure nitric oxide released from endothelial cells in vitro. *Biological Proceeding Online*, 2003; 5: 136-142.
 29. Suzuki, N., Kojima, H., Urano, Y., Kikuchi, K., Hirata, Y., Nagano, T. Orthogonality of calcium concentration and ability of 4, 5 – diamino fluorescein (DAF-2) to detect NO. *Journal of Biology and Chemistry*, 2002; 277: 47- 49.
 30. Wardman, P. Fluorescent and luminescent probes for measurement of oxidative and nitrosative species in cells and tissues: progress, pitfalls, and prospects. *Free Radical Biology and Medicine*, 2007; 43: 995 –1022.
 31. Williams, D.L.H. Nitrosating agents: is peroxynitrite a likely candidate? *Nitric Oxide*, 1997; 1: 522-527.
 32. Wink, D.A., Nims, R.W., Darbyshire, J.F., Christodoulou, D., Hanbauer, I., Cox, G.W., Laval, F., Laval, J., Cook, J.A., Krishna, M.C., DeGraff, W.G., Mitchell, J.B. Reaction kinetics for nitrosation of cysteine and glutathione in aerobic nitric oxide solutions at neutral pH. Insights into the fate and physiological effects of intermediates generated in the NO/O₂ reaction. *Chemical Research and Toxicology*, 1994; 7:519-525.
 33. Zafiriou, O. C. and McFarland, M. Determination of trace levels of nitric oxide in aqueous solution. *Analytical Chemistry*, 1980; 52 (11): 1662–1667.

9/6/2012